

Manual

Clean-Up RNA Concentrator

Kit for RNA preparations concentration and DNA residues removal.

catalog #	size
039-25C	25 isolations
039-100C	100 isolations

For research use only.

Guarantee

A&A Biotechnology provides a guarantee on this product.

The company does not guarantee the correct performance of this kit in the event of:

- not adhering to the supplied protocol
- use of not recommended equipment or materials
- use of other reagents than recommended or which are not a component of the product
- use of expired or improperly stored product or its components



Table of Contents

Contents	3
Additional equipment and reagents	3
Necessary	3
Important notes	3
Purification protocol	
Safety information	6

Contents

component	25 isolations	100 isolations	storage
Microcolumns for RNA purification	25 pcs	100 pcs	15-25 °C
1.5 ml elution tubes	25 pcs	100 pcs	15-25 °C
2 ml tubes	25 pcs	100 pcs	15-25 °C
DNAse, RNAse free (10 U/µI)	60 µl	220 µl	-20 °C
10x DNAse reaction buffer	1.5 ml	1.5 ml	-20 °C
A1 wash solution	30 ml	110 ml	15-25 °C
B1 DNAse removing buffer	20 ml	70 ml	15-25 °C
Ultrapure water	8 ml	15 ml	15-25 °C

The binding capacity of the RNA purification column is 10 μg of RNA.

Additional equipment and reagents

Necessary

- 1.5 ml sterile Eppendorf tubes
- Microcentrifuge
- Heatblock or incubator set to 37 °C

Important notes

When working with RNA, use RNAse-free consumables. Work sterile, use disposable gloves and change them whenever good laboratory practice requires it.

Purification protocol

1.	To 50-100 µl of isolated RNA samples add:
	12 μl of 10x DNAse reaction buffer, 2 μl of DNAse, ultrapure water up to 120 μl.
	Gently mix by pipetting.
2.	Incubate the sample for 15 min at 37 °C .
3.	Add 600 μl of B1 buffer.
4.	Thoroughly mix and apply onto the microcolumn. Close the tube with the cap.
5.	Centrifuge for 30-60 s at 10 000 RPM .
6.	Transfer the microcolumn to a new 2 ml tube (included).
7.	Add 600 µl of A1 wash solution. Close the tube with the cap.
8.	Centrifuge for 1 min at 10 000 RPM .
9.	Remove the microcolumn from the tube and discard the filtrate. Place the microcolumn into the same tube.
10.	Add 400 μl of A1 wash solution. Close the tube with the cap.
11.	Centrifuge for 2 min at 10 000 RPM .
12.	Transfer the microcolumn to a new 1 .5 ml tube (included).
13.	Add $15-30 \mu$ I of ultrapure water directly onto the microcolumn resin. Close the tube with the cap.
14.	Keep for 2 min at room temp .

15. Centrifuge for **1 min** at **10 000 RPM**.

16. Remove the microcolumn and store the tube with purified RNA at -20 °C, -80 °C until later use.

Elution tube has a long, elastic cap connector. Start closing the tube by careful pressing the cap on the connector side. An opening "click" sound confirms proper closure. Different ways of closing may cause opening of the tube during storage.

Safety information

	A1 wash solution
DANGER	H225 Highly flammable liquid and vapor. H319 Causes serious eye irritation. H336 May cause drowsiness or dizziness. P210 Keep away from heat, sparks, open flames, hot surfaces. No smoking. P261 Avoid breathing vapors. P305+P351+P338 If in eyes: rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
• •	B1 DNAse removing buffer
DANGER	H225 Highly flammable liquid and vapor. H319 Causes serious eye irritation. H336 May cause drowsiness or dizziness. P210 Keep away from heat, sparks, open flames, hot surfaces. No smoking. P261 Avoid breathing vapors. P305+P351+P338 If in eyes: rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Manual - Clean-Up RNA Concentrator



A&A Biotechnology, ul. Strzelca 40, 80-299 Gdańsk, Poland phone +48 883 323 761, +48 600 776 268 info@aabiot.com, www.aabiot.com

