

Manual

Sensitive RT HS-PCR Mix SYBR®

High specificity ready-to-use mix for real-time hot-start PCR with SYBR® Green.
2x concentrated.

catalog #	size
2017-100BM	200 reactions in 25 µl
2017-1000BM	2000 reactions in 25 µl

For research use only.

Guarantee

A&A Biotechnology provides guarantee on this product.

The company does not guarantee correct performance of this kit in the event of:

- not adhering to the supplied protocol
- use of not recommended equipment or materials
- use of other reagents than recommended or which are not a component of the product
- use of expired or improperly stored product or its components

SYBR® is a registered trademark of Molecular Probes Inc.

Description

Sensitive RT HS-PCR Mix SYBR[®] is optimized for high specificity ready to use real-time hot-start PCR mixture with SYBR[®] Green.

Mixture contains all components required for qPCR except DNA template and primers.

Taq DNA polymerase is blocked by monoclonal antibody.

The premix formulation saves time and reduces contamination due to a reduced number of pipetting steps required for PCR set up. The mix is optimized for efficient and reproducible reaction.

Contents

	2017-100BM	2017-1000BM	storage
Sensitive SYBR [®]	2 x 1.25 ml	20 x 1.25 ml	-20 °C, in darkness
ultrapure water	2 x 1.5 ml	20 x 1.5 ml	-20 °C

Sensitive SYBR[®] composition

component	amount
modified <i>Taq</i> DNA polymerase	0.1 U/μl
MgCl ₂	4 mM
dNTPs	0.5 mM of each dNTP
2x reaction buffer with SYBR [®] Green	

Notes

- Before use all solutions should be thawed thoroughly on ice, gently mixed by inverting the tube and briefly centrifuged.
- Up to 3x repeated freeze-thaw cycles do not influence the activity of this product.

Example PCR protocol

1.

Thaw **all components of the kit** on ice, gently mix by inverting the tubes and briefly centrifuge. Place the tubes on ice again.
2.

Place PCR tubes on ice and add:

component	PCR reaction volume		
	10 µl	25 µl	50 µl
Sensitive SYBR®	5 µl	12.5 µl	25 µl
primer 1**	0.1-1 µM*	0.1-1 µM*	0.1-1 µM*
primer 2**	0.1-1 µM*	0.1-1 µM*	0.1-1 µM*
DNA, cDNA template	10 pg-1 µg	10 pg-1 µg	10 pg-1 µg
ultrapure water	up to 10 µl	up to 25 µl	up to 50 µl

* recommended for standard qPCR
** final concentration in reaction mixture

3.

Gently vortex the samples and briefly centrifuge to collect all droplets remaining on the tube walls and caps to the bottom of the tube.
4.

Place the tubes in the thermocycler and start the PCR programme.

An example amplification profile:

step	temperature	time
initial denaturation	95 °C	3-5 min
25-45 cycles	95 °C	15-30 s
	50-68 °C	30-60 s
	72 °C	15-60 s*

* depending on the length of PCR products
PCR product melting analysis is recommended.

Recommended ROX mixture

HiROX (0.6-1 µl per 50 µl of total reaction volume): Applied Biosystems: 7000, 7300, 7700, 7900HT Fast, StepOne, StepOnePlus.

LowROX (0.6-1 µl per 50 µl of total reaction volume): Applied Biosystems: 7500, Stratagene: Mx3000P, Mx3005P, Mx4000P.



A&A BIOTECHNOLOGY

innovating life science

A&A Biotechnology, ul. Strzelca 40, 80-299 Gdańsk, Poland
phone +48 883 323 761, +48 600 776 268
info@aabiotech.com, www.aabiotech.com

