

Manual

MagnifiQ™ 1 Plant DNA instant kit

Kit for automated, magnetic isolation of genomic DNA from plant tissue in the strip format. Contains ready-to-use, reagent-filled stripes and all necessary consumables. The strip format enables the isolation of a single sample per purification run.

catalog#	size	compatible devices *
650A-1V-32	32 isolations	Auto-Pure S32 Auto-Pure Mini
650A-1V-160	160 isolations	Auto-Pure S32 Auto-Pure Mini

* Compatible devices

The kit has been tested with specific Allsheng brand isolation devices. This does not preclude it from working with other devices. If your device is not listed, please contact us at info@aabiot.com.

For research use only.

Guarantee

A&A Biotechnology provides a guarantee on this product.

The company does not guarantee the correct performance of this kit in the event of:

- not adhering to the supplied protocol
- use of not recommended equipment or materials
- use of other reagents than recommended or which are not a component of the product
- · use of expired or improperly stored product or its components

Table of Contents

Advantages	4
Specification	4
Description	4
Contents	5
Additional equipment and reagents	5
Necessary	5
Optional	5
Material preparation	6
Dry, powdered plant tissue: 20 - 50 mg	6
Fresh or frozen, powdered plant tissue: 40 - 80 mg	6
Dry, unpowdered plant tissue: 20 - 50 mg	7
Fresh or frozen, unpowdered plant tissue: 40 - 80 mg	7
Protocol	8
Protocol files	8
Extraction protocol	9
Safety information	10

Advantages

- MagnifiQ™ 1 Plant DNA instant kit does not require initial preparation of buffers. Just add samples to the strip and get extracted material within approximately half an hour.
- It enables isolation of different samples with universal kit and automated extraction programme.

Specification

protocol time	~31 min	
sample size	 dry, powdered plant tissue: 20 - 50 mg fresh or frozen, powdered plant tissue: 40 - 80 mg dry, unpowdered plant tissue: 20 - 50 mg fresh or frozen, unpowdered plant tissue: 40 - 80 mg 	
elution volume	100 μΙ	
elution solution	Tris buffer	
binding capacity	up to 60 ug	
downstream applications	sequencing, qPCR, RT-PCR	

Description

MagnifiQ[™] 1 Plant DNA instant kit is designed for genomic DNA isolation from plant tissue.

The isolated material is perfect for further analyzes and tests by PCR and real-time PCR methods and for sequencing.

The MagnifiQ™ product series is based on the automated isolation of nucleic acids with use of magnetic beads. This method significantly shortens working time and reduces risk of mistake in comparison to manual methods.

Contents

650A-1V-32

650A-1V-160

component	quantity	cat#	quantity	cat#	storage
XS-PT - extraction strip	4 x 8 pcs	K-S1V22XPT	20 x 8 pcs	K-S1V22XPT	15-25 ℃
LPE lysis buffer	28 ml	K-LPE-28	140 ml	K-LPE-140	15-25 ℃
Proteinase K	1.1 ml	K-PRK-11A	3 x 1.1 ml	K-PRK-11A	4-8 °C*
L3P precipitation solution	4 ml	K-L3P-4	18 ml	K-L3P-18	15-25 ℃
tip comb 8	8 x 2 pcs	K-C8U-2	40 x 2 pcs	K-C8U-2	15-25 ℃

^{*} Proteinase K can be stored at 15-25 °C for up to 12 months.

Additional equipment and reagents

Necessary

- 2 ml sterile Eppendorf tubes (sample lysis)
- automated pipette
- pipette tips
- Beadbeater
- centrifuge
- thermoblock

Optional

- RNAse (10 μl per sample), cat # 1006-10
- bead-beater tubes M45 (metallic beads), cat # K-2M-25

Material preparation

Dry, powdered plant tissue: 20 - 50 mg


Fresh or frozen, powdered plant tissue: 40 - 80 mg

1. Transfer appropriate amount of powdered plant material to a 2 ml sterile Eppendorf tube (not included). 2. Add 800 µl of LPE buffer and 20 µl Proteinase K. 3. Vortex the sample for 10 s and incubate for 10 min at 50 °C with shaking at 1200 RPM. Centrifuge for 5 min at 14 000 RPM. 4. 5. Transfer 500 µl of supernatant to a 2 ml Eppendorf tube (not included). Optional RNA removal. Add 10 µl of RNAse (cat # 1006-10). Vortex the sample for 10 s and Incubate for 10 min at 50 °C with shaking at 800 RPM. 6. Add 100 µl of L3P precipitation solution and by inverting the tube three times. 7. Put on ice for 3 min. 8. Centrifuge for 10 min at 14 000 RPM. 9. Attention. In the isolation protocol, use $400 \,\mu\text{I}$ of supernatant as the sample. Follow point 1. of the protocol.

Dry, unpowdered plant tissue: 20 - 50 mg Fresh or frozen, unpowdered plant tissue: 40 - 80 mg

Additional reagents you will need:

• bead-beater tubes M45 (metallic beads), cat # K-2M-50




Protocol

Protocol files

device	protocol name	protocol file	installation
Auto-Pure Mini MQ		aabiot.com/protocols/magnifiq/MI/M Q-PLT-MI.txt	Create folder "items" on a USB drive and copy the protocol file to it.
	MQ-PLT-MI		Insert the USB drive into a USB slot in the device.
			 On a device screen, go to Settings > System > Transfer > Import.
			4. Select the protocol and tap "Import".
Auto-Pure Mini (QR code)	MQ-PLT-MI		 On a device screen, go to Run > \(\begin{align*} \begin{align*} \chi_2 \\ \cdot \end{align*} \) Scan the QR code with the device's scanner.
Auto-Pure S32	MQ_PLT_S32	aabiot.com/protocols/magnifig/S32/ MO PLT S32.txt	Create folder "im_export_protocols" on a USB drive and copy the protocol file to it.
			Insert the USB drive into a USB slot in the device.
			3. On a device screen, go to Protocols >Import.
			4. Select the protocol and tap "Import".

Extraction protocol

1. Place **XS-PT** stripes in the rack.



2. Remove the foil from the **XS-PT** stipes starting from well **6**.

Note. The wells are numbered on the side of the strip.

- 3. Add 400 µl of sample to the well 1 (first from the left) on the XS-PT strip.
- 4. Place the rack in the extraction device.
- 5. Place the appropriate number of **tip combs 8** in the extraction device.
- 6. Run the protocol on your device.
- After the program is over, remove the combs and then remove the rack from the extraction device and transfer the purified DNA located in well 6 (first from the right) on the XS-PT strip into sterile tubes (not included).

Note. Store extracted material at 4 °C.

Safety information

DANGER

Proteinase K

H315 Causes skin irritation.

H319 Causes serious eve irritation.

H334 May cause allergy or asthma symptoms or breathing difficulties if inhaled.

H335 May cause respiratory irritation.

P261 Avoid breathing dust.

P305+P351+P338 If in eyes: rinse cautiously with water for several minutes. Remove contact lenses,

if present and easy to do. Continue rinsing.

P342+P311 If experiencing respiratory symptoms call a Poison Center or doctor/physician.

XS-PT - extraction strip















DANGER

H225 Highly flammable liquid and vapor.

H302+H312+H332 Harmful if swallowed, in contact with skin or if inhaled.

H314 Causes severe skin burns and eye damage.

H412 Harmful to aquatic life with long lasting effects.

P210 Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking. P273 Avoid release to the environment.

P280 Wear protective gloves/protective clothing/eye protection/face protection/hearing protection.

P301+P312+P330 If swallowed: Call a poison center/doctor/ if you feel unwell.

P303+P361+P353 If on skin (or hair): Take off immediately all contaminated clothing. Rinse skin with water

 $P304 + P340\ If\ inhaled: Remove\ person\ to\ fresh\ air\ and\ keep\ comfortable\ for\ breathing.$

P305+P351+P338 If in eyes: rinse cautiously with water for several minutes. Remove contact lenses,

if present and easy to do. Continue rinsing.



A&A Biotechnology, ul. Strzelca 40, 80-299 Gdańsk, Poland phone +48 883 323 761, +48 600 776 268 info@aabiot.com, www.aabiot.com

version 2024-3

